


# TRPV4 Antagonism Prevents Mechanically Induced Myotonia

Chris Dupont, BS,<sup>1†</sup> Kevin Novak, PhD,<sup>1†</sup> Kirsten Denman, BS,<sup>1</sup> Jessica H. Myers, BS,<sup>1</sup>  
 Jeremy M. Sullivan, PhD,<sup>2</sup> Phillip V. Walker II BS,<sup>1</sup> Nicklaus L. Brown, BS,<sup>1</sup>  
 David R. Ladle, PhD,<sup>1</sup> Laurent Bogdanik, PhD,<sup>3</sup> Cathleen M. Lutz, PhD,<sup>3</sup>  
 Andrew A. Voss, PhD,<sup>4</sup> Charlotte J. Sumner, MD,<sup>2,5</sup> and Mark M. Rich, MD, PhD <sup>1</sup>

**Objective:** Myotonia is caused by involuntary firing of skeletal muscle action potentials and causes debilitating stiffness. Current treatments are insufficiently efficacious and associated with side effects. Myotonia can be triggered by voluntary movement (electrically induced myotonia) or percussion (mechanically induced myotonia). Whether distinct molecular mechanisms underlie these triggers is unknown. Our goal was to identify ion channels involved in mechanically induced myotonia and to evaluate block of the channels involved as a novel approach to therapy.

**Methods:** We developed a novel system to enable study of mechanically induced myotonia using both genetic and pharmacologic mouse models of myotonia congenita. We extended *ex vivo* studies of excitability to *in vivo* studies of muscle stiffness.

**Results:** As previous work suggests activation of transient receptor potential vanilloid 4 (TRPV4) channels by mechanical stimuli in muscle, we examined the role of this cation channel. Mechanically induced myotonia was markedly suppressed in TRPV4-null muscles and in muscles treated with TRPV4 small molecule antagonists. The suppression of mechanically induced myotonia occurred without altering intrinsic muscle excitability, such that myotonia triggered by firing of action potentials (electrically induced myotonia) was unaffected. When injected intraperitoneally, TRPV4 antagonists lessened the severity of myotonia *in vivo* by approximately 80%.

**Interpretation:** These data demonstrate that there are distinct molecular mechanisms triggering electrically induced and mechanically induced myotonia. Our data indicates that activation of TRPV4 during muscle contraction plays an important role in triggering myotonia *in vivo*. Elimination of mechanically induced myotonia by TRPV4 inhibition offers a new approach to treating myotonia.

ANN NEUROL 2020;88:297–308

Muscle stiffness due to involuntary contraction of muscle (myotonia) is present in a number of dystrophic and nondystrophic muscle diseases.<sup>1–4</sup> The inability of muscle to relax rapidly following cessation of voluntary contraction leads to motor disability, which adversely affects quality of life.<sup>2,4,5</sup> The mainstay of therapy for myotonia is drugs that block Na<sup>+</sup> channels, such as mexiletine.<sup>2,5–8</sup> Block of Na<sup>+</sup> channels reduces muscle excitability such that it is harder to trigger action potentials. Although this may reduce myotonia, it can also lead

to weakness due to impaired action potential generation during voluntary movement.<sup>9</sup> In addition, block of Na<sup>+</sup> channels can alter cardiac excitability increasing the risk of arrhythmias. Many patients incur side effects or incomplete symptom resolution, highlighting the need for novel treatments.<sup>2,5–8</sup>

Myotonia can be triggered in 2 ways: by previous firing of voluntary action potentials or by mechanical stimulation of muscle. Myotonia triggered by voluntary action potentials we term electrically induced myotonia.

View this article online at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1002/ana.25780). DOI: 10.1002/ana.25780

Received Jan 13, 2020, and in revised form Apr 27, 2020. Accepted for publication May 7, 2020.

Address correspondence to Dr Mark Rich, Department of Neuroscience, Cell Biology and Physiology, Wright State University, 3640 Colonel Glenn Highway, Dayton, OH, 45435. E-mail: [mark.rich@wright.edu](mailto:mark.rich@wright.edu)

<sup>†</sup>These authors contributed equally to this work.

From the <sup>1</sup>Department of Neuroscience, Cell Biology and Physiology, Wright State University, Dayton, OH, USA; <sup>2</sup>Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA; <sup>3</sup>The Jackson Laboratory, Bar Harbor, ME, USA; <sup>4</sup>Department of Biology, Wright State University, Dayton, OH, USA; and <sup>5</sup>Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD, USA

Contributors to the generation of electrically induced myotonia include build-up of  $K^+$  in the transverse tubules (t-tubules) and activation of an  $Na^+$  persistent inward current (NaPIC).<sup>10–13</sup> Mechanically induced myotonia can be elicited by tapping muscle with a reflex hammer (percussion myotonia).<sup>1,2</sup> The contribution of mechanically induced myotonia to motor dysfunction has never been determined.

To our knowledge, mechanisms underlying mechanically induced myotonia have not been previously examined. One potential mechanism is the activation of mechanosensitive ion channels. A number of mechanosensitive channels have recently been described and they play central roles in diverse processes, such as the sensation of touch, hearing, and regulation of blood pressure.<sup>14</sup> Skeletal muscle has been shown to express several mechanosensitive channels.<sup>15,16</sup>

Transient receptor potential vanilloid 4 (TRPV4) is a member of the transient receptor potential (TRP) ion channel family that is activated by mechanical stimuli, heat, hypo-osmolarity, and arachidonic acid metabolites.<sup>17,18</sup> In urothelial cells, TRPV4 mediates mechanically evoked elevations of intracellular  $Ca^{2+}$  and plays a central role in regulation of bladder function.<sup>19,20</sup> TRPV4 is expressed in skeletal muscle<sup>21–23</sup> and has been implicated in modulation of resting  $Ca^{2+}$  influx and muscle fatigue.<sup>24,25</sup> Mechanically activated currents present in wild type skeletal muscle are absent in TRPV4-null mice, strongly suggesting that TRPV4 contributes to mechanically activated currents in skeletal muscle.<sup>26</sup>

TRPV4 is most permeant to  $Ca^{2+}$ ,<sup>27</sup> such that its activation would be expected to depolarize muscle. Combining this with the finding that TRPV4 is activated following mechanical stimuli led us to explore the possibility that activation of TRPV4 triggers mechanically induced myotonia.


Our data demonstrate that mechanically induced activation of TRPV4 is a significant contributor to triggering of myotonia in vivo, such that TRPV4 antagonism may offer a new approach to treatment of myotonia.

## Methods

### Mice

All animal procedures were performed in accordance with the policies of the Animal Care and Use Committee of Wright State University. The genetic mouse model of myotonia congenita used was *Cln1<sup>adr-mto</sup>/J* (CIC<sup>adr</sup>) mice, which have a null mutation in the *Cln1* gene (Jackson Laboratory Stock #000939). Genotyping was performed as previously described to select heterozygous mice for breeding.<sup>28</sup> Otherwise, homozygous myotonic mice were identified by appearance and behavior as previously described.<sup>29</sup> To pharmacologically induce myotonia, muscle from asymptomatic littermates was treated with 100  $\mu$ M 9-anthracenecarboxylic acid (9AC) to block CIC-1 chloride channels.<sup>30</sup> Mice were used from 6 weeks to 4 months of age.

TRPV4-null (*Trpv4<sup>-/-</sup>*) mice were generated by homologous recombination using a targeting vector including loxP sites flanking *Trpv4* exons 4 and 5, followed by outcrossing to *Sox2<sup>Cre</sup>*-expressing mice (Jackson Laboratory Stock #004783; Fig 1A). Quantitative reverse transcription polymerase chain reaction (qRT-PCR) analyses confirm the absence of *Trpv4* mRNA expression in these mice (Fig 1B). *Trpv4<sup>-/-</sup>* mice and wild type littermates were generated by interbreeding of *Trpv4<sup>+/-</sup>* mice. Genotyping was performed by polymerase chain reaction (PCR) analysis of tail DNA using the following primers: F1, 5'-GACAGCTGTG TCTCCACCAA; F2, 5'-GTGAGTAGCGGTGGAGG CTA; R1, 5'-GGACAAAATGAGGGTGTGAACTTT



**FIGURE 1:** Generation and characterization of *Trpv4<sup>-/-</sup>* mice. (A) Schematic representation of the targeting strategy. The structure of the endogenous mouse *Trpv4* allele is shown at the top, including the locations of the three genotyping primers, above the targeting vector in which *Trpv4* exons 4 and 5 are flanked by loxP sites. Following excision of the Neo cassette, mice were outcrossed to a *Sox2<sup>Cre</sup>*-expressing mouse strain enabling Cre recombinase (Cre)-mediated gene deletion. (B) *Trpv4* mRNA levels in extensor digitorum longus muscle of adult wild type (WT) and *Trpv4<sup>-/-</sup>* mice, as assessed by quantitative reverse transcription polymerase chain reaction (qRT-PCR; n = 3 for WT and *Trpv4<sup>-/-</sup>*).

(Fig 1A). This multiplex PCR produces a 183-bp amplicon for the wild type allele, a 200-bp amplicon for the null allele, and both amplicons in heterozygous mice. Comparable to previously reported *Trpv4* knockout mouse lines,<sup>31</sup> *Trpv4*<sup>+/-</sup> mice from this newly generated line did not exhibit deficits in weight, survival, or motor function (data not shown).

### Intracellular Recording from Individual Myofibers

Mice were euthanized using CO<sub>2</sub> inhalation followed by cervical dislocation, and both extensor digitorum longus muscles were dissected out tendon-to-tendon. Muscles were maintained at 20 to 22°C and recorded within 6 hours of euthanization. The recording chamber was continuously perfused with Ringer solution containing (in mM): NaCl, 118; KCl, 3.5; CaCl<sub>2</sub>, 1.5; MgSO<sub>4</sub>, 0.7; NaHCO<sub>3</sub>, 26.2; NaH<sub>2</sub>PO<sub>4</sub>, 1.7; and glucose, 5.5 (pH 7.3–7.4), and equilibrated with 95% O<sub>2</sub> and 5% CO<sub>2</sub>.

Intracellular recordings were performed as previously described.<sup>13,29</sup> Briefly, muscles were loaded with 50 μM N-benzyl-p-toluenesulfonamide (BTS) for at least 30 minutes prior to recording to prevent contraction.<sup>32</sup> To aid in visualization of fibers during impalement, muscle was stained for 3 minutes with 10 μM 4-(4-diethylaminostyryl)-N-methylpyridinium iodide (4-Di-2-ASP) and imaged with an upright epifluorescence microscope. Micro-electrodes were filled with 3 M KCl solution containing 1 mM sulforhodamine 101 to visualize the electrodes with epifluorescence. Resistances were between 15 and 30 MΩ, and capacitance compensation was optimized prior to recording. To perform intracellular recordings of action potentials, individual fibers were impaled with 2 sharp electrodes, one to pass current and one to record the voltage response. For comparisons of vehicle treatment to drug treatment, the bilateral extensor digitorum longus (EDL) muscles were dissected out. One was treated with vehicle and the other with drug. This allowed for paired recordings from single mice for vehicle versus drug treatment.

Fibers with resting potentials more depolarized than -74 mV were excluded from analysis. Passive properties and properties of single action potentials were measured as previously described.<sup>28</sup> Action potential voltage threshold was defined as the voltage at which the dV/dt was 0.03 of its maximum (a value near 10 mV/ms) based on methods described by Sekerli et al.<sup>33</sup> Input resistance was calculated by injecting hyperpolarizing current and measuring the steady-state response.

### Quantitative Reverse Transcription PCR

Tissues were homogenized in TRIzol (Thermo Fisher) utilizing Lysing Matrix D 1.4 mm ceramic spheres (MP Biomedicals) and a FastPrep-24 5G Instrument (MP Biomedicals). Total

RNA was then isolated and converted to cDNA as described previously.<sup>34</sup> The qRT-PCR analyses were performed using an ABI 7900HT Real-Time PCR System (Thermo Fisher), with Taqman Universal PCR Master Mix (Thermo Fisher) and the following Taqman Gene Expression Assays: *Gusb*, Mm00446953\_m1 (Thermo Fisher); and *Trpv4*, Mm00499025\_m1 (Thermo Fisher). *Trpv4* transcript levels were quantified using the delta-delta Ct (ΔΔCt) method with normalization to the reference gene *Gusb*. Each reaction was performed in triplicate and all data are presented as average normalized *Trpv4* transcript levels relative to a single wild type sample, which was assigned the value of 1.

### Drugs

HC-067047 was obtained from Hello Bio (Princeton, NJ), GSK2193874 from Tocris (Minneapolis, MN), 9AC and sulforhodamine 101 from Sigma Aldrich (St. Louis, MO), BTS from TCI America (Portland, OR), and 4-Di-2-ASP from Molecular Probes (Eugene, OR).

### In Vivo Force Recording

In vivo muscle force recordings were performed as previously described.<sup>28</sup> Mice were anesthetized via isoflurane inhalation; then the distal tendon of the triceps surae was attached to a force transduction motor and the sciatic nerve stimulated while isometric muscle force generation was measured. Muscle temperature was monitored with a laser probe and maintained between 28°C and 30°C with a heat lamp. The muscle was kept moist by applying mineral oil. Force recordings in *CIC<sup>adr</sup>* mice were performed before and 30 minutes following intraperitoneal (ip) injection. For HC-067047 (20 mg/kg), 0.2 ml of solution containing drug or vehicle was injected (0.015 ml of DMSO and 0.185 ml of saline). GSK2193874 was first dissolved in DMSO at a concentration of 7.5 mg/ml (0.375 mg/50 μl DMSO) using sonication. Distilled water was then added in a 1:1 ratio, followed by another round of sonication. Close to 0.1 ml of this solution was injected ip, with the volume adjusted to administer the correct dose of drug.

### Statistical Analysis and Blinding

For statistical comparisons of intracellular recordings from individual muscle fibers, a repeated measures analysis of variance (ANOVA) with a random effect for mouse was run using SAS version 9.4 (SAS Institute, Cary, NC). The “simulated” post hoc multiple comparison procedure was used for all post hoc multiple comparisons. Origin software (OriginLab, Northampton, MA) was used to perform paired *t*-tests for analysis of in vivo force recordings before and after administration of HC-067047 or GSK2193874. All data are presented as means ± SEM,

$p < 0.05$  was considered to be significant. The numbers of mice and muscle fibers used are described in the corresponding figure legends and text. For statistical analysis “n” was the number of mice.

For ex vivo experiments comparing mechanically induced myotonia in *Trpv4*<sup>+/+</sup> to *Trpv4*<sup>-/-</sup> mice, the experimenter was blinded as to the genotype of the mice. Similarly, ex vivo experiments testing efficacy of drugs were performed with the experimenter blinded as to which vial contained drug and which contained vehicle. For in vivo experiments, as the drug solution was opaque, GSK2193874 solutions were taken up in a syringe by a different experimenter and the syringe was wrapped in tape to maintain blinding until analysis of data was complete.

## Results

We studied muscle isolated from 2 mouse models of myotonia congenita: (1) a pharmacologic model in which ClC-1 chloride channels are blocked with 100  $\mu$ M 9-anthracenecarboxylic acid (the 9AC model)<sup>30</sup>; and (2) a genetic model (*Cln1*<sup>adr-mto</sup>), ClC<sup>adr</sup> mice) in which affected mice are homozygous for a null mutation in the *Cln1* gene.<sup>35</sup> In each model of myotonia, we studied the response of skeletal muscle to both electrical stimulation and mechanical stimulation, 2 distinct triggers of myotonia.

Muscle isolated from wild type mice responded to a 200 ms current injection of depolarizing current by repetitively firing action potentials that immediately ceased when current injection was terminated (Fig 2A). In contrast, muscles from both models of myotonia congenita responded to current injection with firing of action potentials that persisted following cessation of current injection (Fig 2A). We define this as electrically induced myotonia to distinguish it from mechanically induced myotonia.


In order to study mechanically induced myotonia, it was necessary to develop a system that allowed for intracellular recording from individual muscle fibers during mechanical displacement of the fiber. For these studies, fibers were impaled with a single, sharp recording electrode. Approximately 200  $\mu$ m away from the sharp electrode, a blunt 50  $\mu$ m diameter, polished glass pipette was rested against the impaled fiber. The 200  $\mu$ m distance was selected to be as close as possible, yet far enough away that pushing on the fiber with the blunt electrode did not cause loss of impalement (Fig 2B). The blunt electrode was manually advanced approximately 20  $\mu$ m while recording the myofiber's response to mechanical stimulation with the intracellular electrode. This distance was found to trigger mechanically induced myotonia in all

myotonic fibers. To avoid bias in manual advancement of the blunt electrode, the experimenter was blinded as to genotype or drug treatments in all experiments of mechanically induced myotonia.

In wild type muscle, mechanical stimulation triggered a depolarization of  $<3$  mV that did not trigger myotonia (56 fibers from 8 mice; Fig 2C). In contrast, in myofibers from both models of myotonia congenita, 100% of myofibers exhibited mechanically induced myotonia (Fig 2C). To directly measure the depolarization triggered by mechanical stimulation, action potentials were blocked with 1  $\mu$ M tetrodotoxin (TTX; Fig 2D). In myofibers derived from the 9AC model of myotonia congenita, the depolarization averaged  $8.6 \pm 0.5$  mV (35 fibers from 5 mice). In myofibers from the ClC<sup>adr</sup> model, it averaged  $9.5 \pm 0.6$  mV (21 fibers from 3 mice). To our knowledge, this is the first measurement of mechanically induced depolarization of skeletal muscle.

It has been suggested that mechanically activated currents are absent in muscle lacking TRPV4 ion channels.<sup>26</sup> To determine whether TRPV4 might contribute to mechanically induced myotonia, we first assessed *Trpv4* transcript levels in the EDL muscle, the muscle studied physiologically here. The qRT-PCR analyses indicated that *Trpv4* transcript was detectable in the EDL, with equivalent expression levels in wild type and ClC<sup>adr</sup> mice (Fig 3A;  $1.05 \pm 0.07$  wild type vs  $1.19 \pm 0.10$  ClC<sup>adr</sup>;  $p = 0.27$ ).

To determine whether activation of TRPV4 plays a role in triggering mechanically induced myotonia, we recorded from muscle from *Trpv4*<sup>-/-</sup> mice. Muscles from wild type siblings and *Trpv4*<sup>-/-</sup> mice were treated ex vivo with 9AC to cause myotonia and mechanically stimulated with a blunt electrode. In muscles from *Trpv4*<sup>+/+</sup> mice, mechanically induced myotonia was present in 21 of 28 fibers from 4 mice (Fig 3B). In *Trpv4*<sup>-/-</sup> mice, mechanically induced myotonia was present in 0 of 28 fibers from 4 mice (Fig 3A). The elimination of mechanically induced myotonia in 9AC-treated *Trpv4*<sup>-/-</sup> muscle was not accompanied by changes in membrane properties or properties of single action potentials (Table). All 9AC-treated *Trpv4*<sup>+/+</sup> and *Trpv4*<sup>-/-</sup> muscle fibers had electrically induced myotonia (Fig 3C). Following addition of TTX to block action potentials, *Trpv4*<sup>-/-</sup> fibers exhibited a close to 50% reduction in mechanically induced depolarization relative to wild type myofibers ( $5.6 \pm 0.5$  mV in *Trpv4*<sup>+/+</sup> vs  $2.8 \pm 0.5$  in *Trpv4*<sup>-/-</sup>;  $p < 0.01$ ,  $n = 28$  fibers from 4 mice for each group). These data demonstrate that loss of TRPV4 selectively eliminates mechanically induced myotonia by decreasing mechanically induced depolarization of muscle.




**FIGURE 2: Electrically and mechanically induced myotonia are present in both pharmacologic and genetic mouse models of myotonia congenita. (A)** Intracellular recording from muscle fibers during stimulation by injection of depolarizing current. In each of the 3 traces, a 200 ms injection of current (horizontal line under the trace) triggers repeated firing of action potentials during the current injection. In wild type muscle, there is no firing of action potentials following termination of current injection (0/63 fibers from 7 mice). In wild type muscle myofibers in which CIC-1 chloride channels are blocked with 100 μM 9-anthracenecarboxylic acid (9AC) and in muscle from CIC<sup>adr</sup> mice, there is continued firing of action potentials (myotonia) following termination of current injection (58/58 fibers from 8 mice and 72/72 fibers from 9 mice, respectively). **(B)** The arrangement of electrodes used to mechanically stimulate individual muscle fibers. Three muscles fibers are in focus in the image shown. A sharp electrode is impaled into fiber #1 on the right of the image. On the left, a blunt electrode is resting on the bottom of fiber #1, over 200 μm away from the sharp recording electrode. In the top image (no stimulation), the blunt electrode is gently resting on fiber #1. In the lower image, the blunt electrode has been manually advanced to mechanically stimulate fiber #1. **(C)** At the times indicated by vertical arrows, the blunt electrode was manually advanced. In wild type muscle, mechanical stimulation did not trigger myotonia (0/56 fibers from 8 mice). In both 9AC-treated and CIC<sup>adr</sup> muscle, mechanical stimulation triggered myotonia in all fibers tested (42/42 fibers from 6 mice and 56/56 fibers from 8 mice, respectively). **(D)** Examples of mechanically induced depolarization in muscle when 1 μM tetrodotoxin (TTX) was added to the perfusate to block Na<sup>+</sup> channels.

To determine whether small molecule antagonism of TRPV4 might be useful as therapy for myotonia, we examined whether antagonism of TRPV4 could lessen mechanically induced myotonia *ex vivo*. HC-067047 at a dose of 1 μM has previously been shown to fully inhibit TRPV4 in urothelial cells.<sup>20</sup> Muscle from CIC<sup>adr</sup> mice was exposed to either 1 μM HC-067047 or DMSO vehicle at a final concentration of 0.1%. The response to mechanical stimulation was dramatically altered by application of HC-067047. After application of vehicle, 100% of fibers exhibited mechanically induced myotonia (28/28 fibers from 4 mice; Fig 4A), whereas following application of HC-067047, no fibers exhibited mechanically induced myotonia (0/28 fibers from 4 mice; see Fig 4A). To quantitate the effect of HC-067047 on the mechanically induced depolarization, 1 μM TTX was added to the bath to block generation of action potentials. CIC<sup>adr</sup> muscle fibers depolarized by  $9.5 \pm 0.6$  mV following mechanical stimulation. Treatment with 1 μM HC-067047 decreased

this to  $5.5 \pm 0.6$  mV ( $p < 0.01$  vs vehicle;  $n = 21$  vehicle-treated fibers from 3 mice and 21 HC-067047-treated fibers from 3 mice). Application of HC-067047 had no effect on mechanically induced depolarization in *Trpv4*<sup>-/-</sup> muscle ( $2.8 \pm 0.5$  vs  $2.7 \pm 0.4$  mV;  $n = 28$  fibers from 4 mice for each group). These data strongly suggest that HC-067047 abolishes mechanically induced myotonia by blocking TRPV4 channels.

Application of HC-067047 did not eliminate electrically induced myotonia (Fig 4B). After treatment with either vehicle or HC-067047, electrically induced myotonia was present in all fibers (33/33 vehicle-treated fibers from 5 mice and 48/48 HC-067047-treated fibers from 5 mice). Similar results were obtained using 1 μM GSK2193874 to block TRPV4 channels.<sup>36</sup> Application of GSK2193874 greatly reduced the incidence of mechanically induced myotonia (19/20 fibers with mechanically induced myotonia in 3 mice treated with vehicle and 3/20 fibers with mechanically induced myotonia in 3 mice



**FIGURE 3:** Genetic deletion of transient receptor potential vanilloid 4 (TRPV4) channels eliminates mechanically induced myotonia without affecting electrically induced myotonia. (A) Scatter plot of the relative *Trpv4* transcript levels normalized to the reference gene *Gusb* for 6 wild type and 6 *C1cn1<sup>adr-mto</sup>/J* (*C1c<sup>adr</sup>*) extensor digitorum longus muscles. The horizontal bars represent the mean of each group. (B) In 9-anthracenecarboxylic acid (9AC)-treated muscle from *Trpv4<sup>+/+</sup>* mice, mechanically induced myotonia was present in 21 of 28 fibers from 4 mice, whereas in 9AC-treated muscle fibers from *Trpv4<sup>-/-</sup>* mice, mechanically induced myotonia was present in 0 of 28 fibers from 4 mice. At the times indicated by vertical arrows, a blunt electrode was manually advanced to mechanically stimulate the fiber being studied. (C) Electrically induced myotonia was present in 35 of 35 9AC-treated *Trpv4<sup>+/+</sup>* fibers from 4 mice and 22 of 22 *Trpv4<sup>-/-</sup>* fibers from 3 mice.

**TABLE. Genetic Deletion and Pharmacologic Inhibition of TRPV4 Have No Effect on Skeletal Muscle Passive Properties and Single Action Potentials**


	Resting potential, mV	Input resistance, M $\Omega$	Action potential		
			Threshold, mV	Maximum rate of rise, mV/ms	Peak, mV
TRPV4 <sup>+/+</sup> + 9AC	-83.9 $\pm$ 0.4	1.17 $\pm$ 0.09	-62.7 $\pm$ 0.2	239.4 $\pm$ 7.6	32.5 $\pm$ 0.9
TRPV4 <sup>-/-</sup> + 9AC	-83.6 $\pm$ 0.8	1.20 $\pm$ 0.07	-62.9 $\pm$ 0.7	231.9 $\pm$ 2.0	29.6 $\pm$ 0.6
<i>C1c<sup>adr</sup></i> + vehicle	-80.9 $\pm$ 0.5	0.85 $\pm$ 0.03	-56.7 $\pm$ 0.5	234.1 $\pm$ 30.4	30.1 $\pm$ 4.7
<i>C1c<sup>adr</sup></i> + HC-067047	-80.9 $\pm$ 0.5	0.79 $\pm$ 0.05	-56.9 $\pm$ 0.6	236.6 $\pm$ 30.6	29.9 $\pm$ 3.9
<i>C1c<sup>adr</sup></i> + vehicle	-84.5 $\pm$ 0.3	1.06 $\pm$ 0.04	-60.4 $\pm$ 0.4	244.9 $\pm$ 15.0	35.7 $\pm$ 1.0
<i>C1c<sup>adr</sup></i> + GSK2193874	-82.8 $\pm$ 1.4	1.15 $\pm$ 0.07	-59.8 $\pm$ 1.0	242.8 $\pm$ 2.7	36.6 $\pm$ 0.7

For studies of HC-067047, n = 28 fibers from 4 muscles for both groups. For studies of GSK2193874, n = 25 fibers from 3 muscles for vehicle and 19 fibers from 3 muscles for drug. For studies of TRPV4<sup>-/-</sup> mice, n = 39 fibers from 3 TRPV4<sup>+/+</sup> mice, and 22 fibers from 3 TRPV4<sup>-/-</sup> mice. Data shown are mean  $\pm$  SEM. None of the differences between vehicle and drug treatment groups and between TRPV4 genotypes were statistically significant.

9AC = 9-anthracenecarboxylic acid; *C1c<sup>adr</sup>* = *C1cn1<sup>adr-mto</sup>/J*; TRPV4 = transient receptor potential vanilloid 4.

treated GSK2193874; Fig 4C) while having no effect on the incidence of electrically induced myotonia (21/21 fibers from 3 mice for vehicle and 19/19 fibers from 3 mice for GSK2193874; Fig 4D). These data suggest that drugs that block TRPV4 selectively antagonize mechanically induced myotonia.


Muscle excitability is determined by the amount of depolarizing current or depolarization of membrane potential required to trigger an action potential. In hyperexcitable myotonic muscle, muscle is too easily depolarized to action potential threshold, whereas in hypoexcitable muscle it is difficult to reach action potential threshold.



**FIGURE 4:** Treatment with small molecule inhibitors of transient receptor potential vanilloid 4 (TRPV4) greatly reduces mechanically induced myotonia in *Clcn1<sup>adr-mto</sup>/J* (*Clc<sup>adr</sup>*) muscle. (A) Mechanically induced myotonia was present in 28 of 28 vehicle-treated fibers from 4 mice, and 0 of 28 fibers treated with 1  $\mu$ M HC-067047 from 4 mice. At the times indicated by vertical arrows, a blunt electrode was manually advanced to mechanically stimulate the fiber being studied. (B) Electrically induced myotonia was present in 33 of 33 vehicle-treated fibers from 5 mice and 48 of 48 HC-067047-treated fibers from 5 mice. (C) Following application of vehicle 19 of 20 fibers from 3 mice had mechanically induced myotonia. After application of GSK2193874, 3 of 20 fibers from 3 mice had mechanically induced myotonia. (D) Electrically induced myotonia was present in 21 of 21 vehicle-treated fibers from 3 mice and 19 of 19 fibers treated with 1  $\mu$ M GSK2193874 from 3 mice.


To determine whether HC-067047 and GSK2193874 were eliminating mechanically induced myotonia via a reduction in excitability, we measured passive membrane

properties and properties of single action potentials. These properties of muscle determine the difficulty in reaching action potential threshold. Resting potential, input




**FIGURE 5:** Block of transient receptor potential vanilloid 4 (TRPV4) is effective in treating myotonia in vivo. (A) The response of wild type muscle to 2 seconds of 60 Hz stimulation of the sciatic nerve. The horizontal bar under the trace indicates the period of 60 Hz stimulation of the sciatic nerve. Force is fully fused and there is immediate, complete relaxation following termination of stimulation. (B) The response of muscle from *Clcn1<sup>adr-mto</sup>/J* (*Clc<sup>adr</sup>*) mice to 60 Hz stimulation. The vertical dotted line in each trace represents the rapid relaxation of force that occurs in wild type muscle following termination of nerve stimulation. Pre-injection, force generation during 60 Hz stimulation is fused normally, but relaxation following termination of stimulation is only partial and there is continued contraction for many seconds secondary to myotonia. At 45 minutes postinjection of vehicle, myotonia is minimally changed, but is greatly decreased in mice that received either HC-067047 or GSK2193874. (C) Scatter plot of the ratio of severity of myotonia postinjection to severity of myotonia pre-injection in each mouse. The horizontal bars represent the mean of each group. Following injection of vehicle, there was no change in severity of myotonia. Following injection of either drug, there was a marked reduction. \*\*Indicates  $p < 0.01$  versus vehicle.

### Electrically-Induced Myotonia



### Mechanically-Induced Myotonia



**FIGURE 6:** Distinct mechanisms underlying electrically- and mechanically induced myotonia. Electrically induced myotonia (top two rows): in wild type muscle  $\text{Cl}^-$  current plays a central role in controlling resting potential both at rest and during firing of action potentials during voluntary movement. Because of the stabilizing influence of  $\text{Cl}^-$  current,  $\text{K}^+$  build-up in transverse tubules (t-tubules) does not cause sufficient depolarization to trigger myotonia following voluntary activation of muscle. In myotonia congenita, muscle  $\text{Cl}^-$  current is reduced such that  $\text{K}^+$  build-up causes depolarization large enough to trigger myotonia. Mechanically induced myotonia (bottom two rows): mechanical stimulation of wild type muscle activates transient receptor potential vanilloid 4 (TRPV4) channels, which depolarizes muscle. This depolarization is opposed by  $\text{Cl}^-$  current, such that myotonia is not triggered. Activation of TRPV4 channels during mechanical stimulation of muscle with reduced  $\text{Cl}^-$  current causes depolarization sufficient to trigger myotonia.

resistance, and properties of single action potentials were unchanged by either drug (see Table), suggesting neither drug altered excitability.

To determine whether antagonism of TRPV4 might be effective as therapy, we measured the severity of myotonia using in vivo contractile force recordings of gastrocnemius muscle as described previously.<sup>28</sup> The sciatic nerve was stimulated at 60 Hz while measuring isometric muscle force generation. In wild type muscle, force fused during 60 Hz stimulation and relaxed immediately upon cessation of stimulation (Fig 5A). In  $\text{ClC}^{\text{adr}}$  muscle, fusion of force was normal during stimulation, but relaxation following termination of stimulation was only partial due to

myotonia (Fig 5B). Both amplitude and duration of muscle force have been previously used to estimate severity of myotonia.<sup>37</sup> To combine these into a single value, we took the integral of force with respect to time.<sup>9</sup> To eliminate differences in measurement of myotonia due to differences in muscle force between mice, the integral of myotonia was normalized to force generated by a single stimulus before and after administration of drug. We measured severity of myotonia before and 30 minutes after ip injection of 25 mg/kg GSK2193874, 20 mg/kg HC-067047 or vehicle in  $\text{ClC}^{\text{adr}}$  mice. Following injection of vehicle there was no change in severity of myotonia (Fig 5B, C). Following injection of both GSK2193874 and HC-067047



myotonia was reduced by over 80% (Fig 5B, C,  $p < 0.01$  vs vehicle for both drugs,  $n = 5$  mice for each drug treatment group and 6 for vehicle).

## Discussion

Current treatment of myotonia is only partially effective and can cause unwanted side effects due, in part, to non-specific suppression of excitability of both cardiac and skeletal muscle. Identifying currents in skeletal muscle that contribute to triggering of myotonia may lead to development of better targeted therapies that treat muscle stiffness secondary to myotonia without unwanted side effects. We established a novel *ex vivo* system to study currents responsible for mechanically induced (percussion) myotonia. Genetic deletion of TRPV4 suppressed generation of mechanically induced myotonia, while having no effect on myotonia triggered by previous firing of action potentials (electrically induced myotonia). These data provide the first definitive evidence of distinct molecular mechanisms underlying electrically and mechanically induced myotonia. Small molecule inhibitors of TRPV4 reduced the incidence of mechanically induced myotonia *ex vivo* and treatment of mice with systemic administration of these inhibitors also significantly reduced myotonia triggered by repetitive stimulation of the sciatic nerve *in vivo*. Our findings suggest antagonism of TRPV4 may offer a novel therapeutic approach to treating myotonia. A particular advantage of TRPV4 blockade is that it did not affect muscle excitability (the ease of generating action potentials) such that unwanted side effects of weakness and altered cardiac excitability might be avoided.

### Two Molecularly Distinct Triggers of Myotonia

It has been known for many years that myotonia can be triggered by either voluntary firing of action potentials or percussion of muscle.<sup>1,2</sup> We term these electrically and mechanically induced myotonia, respectively. Although it has been presumed that distinct mechanisms are involved, direct proof has been lacking. One reason for the lack of progress in identifying the currents responsible for generation of mechanically induced myotonia has been the inability to record intracellularly from intact muscle fibers during mechanical stimulation of muscle. We developed a system that allowed for intracellular recording of the depolarization of membrane potential and the subsequent myotonia triggered by mechanical stimulation of individual fibers. This made it possible to determine that genetic deletion of TRPV4 ion channels greatly reduced mechanically induced myotonia, while having no effect on electrically induced myotonia.

Shown in Figure 6 is a depiction of our current understanding of the distinct mechanisms underlying electrically and mechanically induced myotonia in the muscle

disease myotonia congenita. In myotonia congenita, there is a reduction in the number of functional  $\text{Cl}^-$  channels.  $\text{ClC-1}$ -mediated  $\text{Cl}^-$  conductance accounts for 70 to 80% of resting muscle membrane conductance and stabilizes the membrane potential near the resting potential.<sup>35,38–40</sup> In the absence of  $\text{Cl}^-$  current, stimuli that would otherwise cause only modest depolarization led to depolarization large enough to trigger myotonia.

In electrically induced myotonia, an important source of depolarization is exit of  $\text{K}^+$  from the fiber during repolarization of action potentials. The exit of  $\text{K}^+$  leads to build-up of  $\text{K}^+$  in t-tubules, a depolarized shift in the  $\text{K}^+$  equilibrium potential, and depolarization of the membrane potential.<sup>10–12,38</sup> When the  $\text{Cl}^-$  current is small or absent the depolarization caused by  $\text{K}^+$  build-up triggers myotonia.

In mechanically induced myotonia, there have been no preceding action potentials so build-up of  $\text{K}^+$  in t-tubules cannot be a contributor to depolarization. Instead, mechanical distortion of the sarcolemma results in opening of TRPV4 channels, which are nonselectively permeable to monovalent and divalent cations,<sup>27</sup> such that their activation causes depolarization and myotonia. A prediction of this model of mechanically induced myotonia is that agonists of TRPV4 channels would worsen myotonia, although this remains to be tested experimentally.

*Ex vivo* experiments demonstrated that electrically and mechanically induced myotonia could be triggered independently. However, *in vivo*, block of TRPV4 channels significantly reduced the severity of myotonia triggered by repetitive stimulation of the sciatic nerve (electrically induced myotonia). The difference is likely due to experimental conditions. During *ex vivo* studies muscle contraction was eliminated with BTS, a drug that blocks interaction of actin and myosin<sup>32,41</sup> because contraction of muscle during intracellular recording causes damage-induced loss of membrane potential. During *in vivo* experiments, muscle contraction was not blocked so that force could be measured. Thus, firing of action potentials *in vivo* increased tension as muscle contracted, which seems to activate TRPV4 to worsen myotonia. We conclude that build-up of  $\text{K}^+$  in t-tubules and activation of TRPV4 combine to trigger myotonia *in vivo*. The lessening of myotonia following block of TRPV4 strongly suggests mechanically induced activation of TRPV4 is a contributor to generation of myotonia *in vivo*.

### Mechanisms Underlying Mechanically-Induced Depolarization of Muscle

The simplest explanation for the contribution of TRPV4 to mechanically induced myotonia is that TRPV4 is activated by mechanical forces exerted on the sarcolemma. Several *in vitro* studies have provided evidence that TRPV4 is

insensitive to membrane stretch,<sup>42-45</sup> but can be activated by mechanical stimuli applied at cell-substrate contacts<sup>43</sup> or to cell surface  $\beta$ 1-integrins.<sup>46</sup> Given the importance of cell-matrix interactions to skeletal muscle physiology,<sup>47</sup> cell-substrate contacts may play a key role in sarcolemma TRPV4 activation in response to mechanical stimulation. However, little is currently known about how TRPV4 is activated by mechanical stimuli in vivo. It also remains to be determined whether the biomechanical properties of skeletal muscle change in the context of myotonia and whether such changes alter the sensitivity of TRPV4 to mechanical stimuli.

In several cell types TRPV4 activation in response to osmomechanical stress occurs with a latency of seconds and is mediated by phospholipase A2 (PLA<sub>2</sub>)-dependent generation of second messengers, such as arachidonic acid.<sup>48-51</sup> In contrast, TRPV4 activation by deflection at cell-substrate interfaces occurs with a rapid latency (<2 ms) and is PLA<sub>2</sub>-independent, consistent with direct transduction of the mechanical stimulus.<sup>43</sup> Similarly, calcium influx through TRPV4 resulting from the application of mechanical force to  $\beta$ 1-integrins occurs with a latency of milliseconds.<sup>46</sup> The rapid onset of mechanically induced myotonia observed in both the pharmacologic and genetic models described here (see Fig 2) suggests TRPV4 activation in this context may be mediated by mechanisms comparable to those underlying channel responses to cell-substrate deflections or integrin stimulation, rather than via activation by second messengers.

It remains to be determined whether all mechanically activated currents contributing to triggering of myotonia have been identified. Although there is one study suggesting genetic deletion of TRPV4 abrogates mechanically activated currents in muscle,<sup>26</sup> we did not observe elimination of mechanically induced depolarization. The remaining depolarization may be due to a technical issue, such as depolarization by the impaling electrode during mechanical stimuli. Alternatively, a second mechanically activated channel expressed in muscle, such as Piezo1, TRPC1, TRPC6, or TRPV2, might also contribute to generation of mechanically induced myotonia.<sup>21,22,52-55</sup>

### TRPV4 Antagonists as a Novel Approach to Therapy of Myotonia

Myotonia is present in muscle diseases, such as myotonia congenita, myotonic dystrophy, sodium channel myotonia, paramyotonia congenita, and hyperkalemic periodic paralysis.<sup>1-4</sup> Myotonia causes functionally limiting muscle stiffness, weakness, and pain.<sup>2,4,5</sup> For movement to occur normally, muscles need to sequentially contract and relax during flexion and extension of a joint. For this to occur, the muscle must fire action potentials during contraction, and be electrically silent during relaxation, as antagonist muscles contract and the muscle is stretched as it lengthens. If muscle is unable to relax

while lengthening, due to mechanically induced myotonia, this will negatively impact motor performance. The contribution of mechanically induced myotonia to motor dysfunction in patients with myotonia has never been determined, as it has not been possible to selectively block mechanically induced myotonia. As described above, our studies suggest mechanically induced myotonia is a significant contributor to motor dysfunction in vivo, such that block of mechanically induced myotonia may improve motor performance.

The current approach to therapy of myotonia is focused on the block of Na<sup>+</sup> channels responsible for generation of action potentials.<sup>2,5-8</sup> This reduces excitability and, thus, reduces myotonia. A problem with this approach is that activation of Na<sup>+</sup> channels is required during the action potentials necessary for voluntary movement. Thus, care must be taken to not block too high a percentage of Na<sup>+</sup> channels, so as not to cause weakness/paralysis.

A superior approach to therapy would be to block pathologic excitation of muscle without affecting normal excitation.<sup>9</sup> Knockout of TRPV4 had no effect on properties of single action potentials that depend on Na<sup>+</sup> current, suggesting that excitability is not significantly reduced. Thus, it may be possible to reduce myotonia by blocking TRPV4 without adversely affecting strength during voluntary movement. It has been reported that there are muscle transcriptional changes in *Trpv4*<sup>-/-</sup> mice, and that TRPV4 activation modulates resting Ca<sup>2+</sup> influx and muscle fatigue,<sup>24,25</sup> such that the block of TRPV4 may have side effects related to muscle function. However, despite TRPV4 being expressed in a number of tissues, knockout mice have a surprisingly mild phenotype, suggesting that the block of TRPV4 in vivo might not lead to intolerable side effects.<sup>31</sup> Administration of selective TRPV4 blockers is well tolerated in both rodents and patients.<sup>36,56</sup> Thus, using the block of TRPV4 as therapy for patients with myotonia may become possible in the near future. One approach would be to add block of TRPV4 to current therapy, as there would be little risk of causing muscle hypoexcitability.

The studies reported here focused on myotonia resulting from loss of muscle Cl<sup>-</sup> current. However, as TRPV4 channels are present in wild type muscle, it is possible that activation of TRPV4 contributes to muscle stiffness in other diseases with myotonia. Thus, the block of TRPV4 may be useful in treating myotonia in diseases such as myotonic dystrophy, sodium channel myotonia, paramyotonia congenita, and hyperkalemic periodic paralysis.

### Acknowledgments

This work was supported by NIH grants AR074985 (M.M.R.), NS087579 (C.J.S), and NS099850 (A.A.V.), and MDA grant 602459 (M.M.R.).

## Author Contributions

M.R., C.S., and A.V. contributed to the conception and design of the study. C.D., K.N., K.D., J.M., J.S., P.W., N.B., D.L., L.B., and C.L. contributed to acquisition and analysis of data. C.D., K.N., K.D., J.M., and J.S. contributed to preparing the figures. J.S., A.V., C.S., and M.R. drafted the manuscript.

## Potential Conflicts of Interest

The authors declared no conflict of interest.

## References

- Trivedi JR, Bundy B, Statland J, et al. Non-dystrophic myotonia: prospective study of objective and patient reported outcomes. *Brain* 2013;136:2189–2200.
- Cannon SC. Channelopathies of skeletal muscle excitability. *Compr Physiol* 2015;5:761–790.
- Thornton CA. Myotonic dystrophy. *Neurol Clin* 2014;32:705–719, viii.
- Cannon SC. Sodium channelopathies of skeletal muscle. *Handb Exp Pharmacol* 2018;246:309–330.
- Statland JM, Bundy BN, Wang Y, et al. Mexiletine for symptoms and signs of myotonia in nondystrophic myotonia: a randomized controlled trial. *JAMA* 2012;308:1357–1365.
- Lehmann-Horn F, Jurkat-Rott K, Rudel R. Diagnostics and therapy of muscle channelopathies—guidelines of the Ulm Muscle Centre. *Acta Myol* 2008;27:98–113.
- Trivedi JR, Cannon SC, Griggs RC. Nondystrophic myotonia: challenges and future directions. *Exp Neurol* 2014;253:28–30.
- Matthews E, Hanna MG. Repurposing of sodium channel antagonists as potential new anti-myotonic drugs. *Exp Neurol* 2014;261:812–815.
- Metzger S, Dupont C, Voss AA, Rich MM. Central role of subthreshold currents in myotonia. *Ann Neurol* 2020;87:175–183.
- Adrian RH, Marshall MW. Action potentials reconstructed in normal and myotonic muscle fibres. *J Physiol* 1976;258:125–143.
- Wallinga W, Meijer SL, Alberink MJ, et al. Modelling action potentials and membrane currents of mammalian skeletal muscle fibres in coherence with potassium concentration changes in the T-tubular system. *Eur Biophys J* 1999;28:317–329.
- Fraser JA, Huang CL, Pedersen TH. Relationships between resting conductances, excitability, and t-system ionic homeostasis in skeletal muscle. *J Gen Physiol* 2011;138:95–116.
- Hawash AA, Voss AA, Rich MM. Inhibiting persistent inward sodium currents prevents myotonia. *Ann Neurol* 2017;82:385–395.
- Ranade SS, Syeda R, Patapoutian A. Mechanically activated ion channels. *Neuron* 2015;87:1162–1179.
- Franco-Obregon A, Lansman JB. Changes in mechanosensitive channel gating following mechanical stimulation in skeletal muscle myotubes from the mdx mouse. *J Physiol* 2002;539:391–407.
- Yeung EW, Whitehead NP, Suchyna TM, et al. Effects of stretch-activated channel blockers on  $[Ca^{2+}]_i$  and muscle damage in the mdx mouse. *J Physiol* 2005;562:367–380.
- Darby WG, Grace MS, Baratchi S, McIntyre P. Modulation of TRPV4 by diverse mechanisms. *Int J Biochem Cell Biol* 2016;78:217–228.
- Shibasaki K. TRPV4 ion channel as important cell sensors. *J Anesth* 2016;30:1014–1019.
- Mochizuki T, Sokabe T, Araki I, et al. The TRPV4 cation channel mediates stretch-evoked  $Ca^{2+}$  influx and ATP release in primary urothelial cell cultures. *J Biol Chem* 2009;284:21257–21264.
- Everaerts W, Zhen X, Ghosh D, et al. Inhibition of the cation channel TRPV4 improves bladder function in mice and rats with cyclophosphamide induced cystitis. *Proc Natl Acad Sci U S A* 2010;107:19084–19089.
- Brinkmeier H. TRP channels in skeletal muscle: gene expression, function and implications for disease. *Adv Exp Med Biol* 2011;704:749–758.
- Gailly P. TRP channels in normal and dystrophic skeletal muscle. *Curr Opin Pharmacol* 2012;12:326–334.
- Blackburn DM, Lazure F, Corchado AH, et al. High-resolution genome-wide expression analysis of single myofibers using SMART-Seq. *J Biol Chem* 2019;294:20097–20108.
- Kusudo T, Wang Z, Mizuno A, et al. TRPV4 deficiency increases skeletal muscle metabolic capacity and resistance against diet induced obesity. *J Appl Physiol* (1985) 2012;112:1223–1232.
- Pritschow BW, Lange T, Kasch J, et al. Functional TRPV4 channels are expressed in mouse skeletal muscle and can modulate resting  $Ca^{2+}$  influx and muscle fatigue. *Pflugers Arch* 2011;461:115–122.
- Ho TC, Horn NA, Huynh T, et al. Evidence TRPV4 contributes to mechanosensitive ion channels in mouse skeletal muscle fibers. *Channels (Austin)* 2012;6:246–254.
- Voets T, Prenen J, Vriens J, et al. Molecular determinants of permeation through the cation channel TRPV4. *J Biol Chem* 2002;277:33704–33710.
- Dupont C, Denman KS, Hawash AA, et al. Treatment of myotonia congenita with retigabine in mice. *Exp Neurol* 2019;315:52–59.
- Novak KR, Norman J, Mitchell JR, et al. Sodium channel slow inactivation as a therapeutic target for myotonia congenita. *Ann Neurol* 2015;77:320–332.
- Palade PT, Barchi RL. On the inhibition of muscle membrane chloride conductance by aromatic carboxylic acids. *J Gen Physiol* 1977;69:879–896.
- Grace MS, Bonvini SJ, Belvisi MG, McIntyre P. Modulation of the TRPV4 ion channel as a therapeutic target for disease. *Pharmacol Ther* 2017;177:9–22.
- Macdonald WA, Pedersen TH, Clausen T, Nielsen OB. N-Benzyl-p-toluene sulphonamide allows the recording of trains of intracellular action potentials from nerve-stimulated intact fast-twitch skeletal muscle of the rat. *Exp Physiol* 2005;90:815–825.
- Sekerli M, Del Negro CA, Lee RH, Butera RJ. Estimating action potential thresholds from neuronal time-series: new metrics and evaluation of methodologies. *IEEE Trans Biomed Eng* 2004;51:1665–1672.
- Ramos DM, d'Ydewalle C, Gabbeta V, et al. Age-dependent SMN expression in disease-relevant tissue and implications for SMA treatment. *J Clin Invest* 2019;129:4817–4831.
- Steinmeyer K, Klocke R, Ortland C, et al. Inactivation of muscle chloride channel by transposon insertion in myotonic mice. *Nature* 1991;354:304–308.
- Thorneloe KS, Cheung M, Bao W, et al. An orally active TRPV4 channel blocker prevents and resolves pulmonary edema induced by heart failure. *Sci Transl Med* 2012;4:159ra48.
- van Lunteren E, Spiegler SE, Moyer M. Fatigue-inducing stimulation resolves myotonia in a drug induced model. *BMC Physiol* 2011;11:5.
- Adrian RH, Bryant SH. On the repetitive discharge in myotonic muscle fibres. *J Physiol* 1974;240:505–515.
- Palade PT, Barchi RL. Characteristics of the chloride conductance in muscle fibers of the rat diaphragm. *J Gen Physiol* 1977;69:325–342.

40. Steinmeyer K, Ortland C, Jentsch TJ. Primary structure and functional expression of a developmentally regulated skeletal muscle chloride channel. *Nature* 1991;354:301–304.
41. Cheung A, Dantzig JA, Hollingworth S, et al. A small-molecule inhibitor of skeletal muscle myosin II. *Nat Cell Biol* 2002;4:83–88.
42. Strotmann R, Harteneck C, Nunnenmacher K, et al. OTRPC4, a non-selective cation channel that confers sensitivity to extracellular osmolarity. *Nat Cell Biol* 2000;2:695–702.
43. Servin-Vences MR, Moroni M, Lewin GR, Poole K. Direct measurement of TRPV4 and PIEZO1 activity reveals multiple mechanotransduction pathways in chondrocytes. *Elife* 2017;30:6.
44. Nikolaev YA, Cox CD, Ridone P, et al. Mammalian TRP ion channels are insensitive to membrane stretch. *J Cell Sci* 2019;132.
45. Martinac B, Poole K. Mechanically activated ion channels. *Int J Biochem Cell Biol* 2018;97:104–107.
46. Matthews BD, Thodeti CK, Tytell JD, et al. Ultra-rapid activation of TRPV4 ion channels by mechanical forces applied to cell surface beta1 integrins. *Integr Biol (Camb)* 2010;2:435–442.
47. Goody MF, Sher RB, Henry CA. Hanging on for the ride: adhesion to the extracellular matrix mediates cellular responses in skeletal muscle morphogenesis and disease. *Dev Biol* 2015;401:75–91.
48. Vriens J, Watanabe H, Janssens A, et al. Cell swelling, heat, and chemical agonists use distinct pathways for the activation of the cation channel TRPV4. *Proc Natl Acad Sci U S A* 2004;101:396–401.
49. Vriens J, Owsianik G, Fisslthaler B, et al. Modulation of the Ca<sup>2+</sup> permeable cation channel TRPV4 by cytochrome P450 epoxygenases in vascular endothelium. *Circ Res* 2005;97:908–915.
50. Ryskamp DA, Jo AO, Frye AM, et al. Swelling and eicosanoid metabolites differentially gate TRPV4 channels in retinal neurons and glia. *J Neurosci* 2014;34:15689–15700.
51. Toft-Bertelsen TL, Yarishkin O, Redmon S, et al. Volume sensing in the transient receptor potential vanilloid 4 ion channel is cell type-specific and mediated by an N-terminal volume-sensing domain. *J Biol Chem* 2019;294:18421–18434.
52. Maroto R, Raso A, Wood TG, et al. TRPC1 forms the stretch-activated cation channel in vertebrate cells. *Nat Cell Biol* 2005;7:179–185.
53. Spassova MA, Hewavitharana T, Xu W, et al. A common mechanism underlies stretch activation and receptor activation of TRPC6 channels. *Proc Natl Acad Sci U S A* 2006;103:16586–16591.
54. Alessandri-Haber N, Dina OA, Chen X, Levine JD. TRPC1 and TRPC6 channels cooperate with TRPV4 to mediate mechanical hyperalgesia and nociceptor sensitization. *J Neurosci* 2009;29:6217–6228.
55. Zanou N, Mondin L, Fuster C, et al. Osmosensation in TRPV2 dominant negative expressing skeletal muscle fibres. *J Physiol* 2015;593:3849–3863.
56. Goyal N, Skrdla P, Schroyer R, et al. Clinical pharmacokinetics, safety, and tolerability of a novel, first-in-class TRPV4 ion channel inhibitor, GSK2798745, in healthy and heart failure subjects. *Am J Cardiovasc Drugs* 2019;19:335–342.